Supplemental Material

Journal of Climate
Seasonal Prediction and Predictability of Regional Antarctic Sea Ice
https://doi.org/10.1175/JCLI-D-20-0965.1

Supplementary Material for "Seasonal prediction and predictability of regional Antarctic sea ice"

Mitchell Bushuk, Michael Winton, Alexander Haumann, Thomas Delworth, Feiyu Lu,Yongfei Zhang, Liwei Jia, Liping Zhang, William Cooke, Matthew Harrison, Bill Hurlin, Nathaniel C. Johnson, Sarah Kapnick, Colleen McHugh, Hiroyuki Murakami, Anthony Rosati, Kai-Chih Tseng, Andrew T. Wittenberg, Xiaosong Yang, Fanrong Zeng

S1 Regional Prediction Skill: ACC



Figure S1: As in Figure 5, but for non-detrended time series.

S2 Regional Prediction Skill: MSSS



Figure S2: As in Figure 6, but for non-detrended time series.

S3 Regional Prediction Skill: ACC^2



Figure S3: As in Figure 6, but for squared detrended ACC values.

S4 Regional Prediction Skill: MSSS



Figure S4: As in Figure 6, but for time series that have been mean-biased corrected by an additive correction, but have not been linear regression adjusted.

S5 Weddell SIE-SIT Correlations in SPEAR_LO



Figure S5: As in Figure 11, but for SPEAR_LO.

S6 Winter sea ice drift climatologies



Figure S6: Climatological JAS sea ice drift patterns from the experiments that produce sea ice ICs in (b) FLOR, (c) SPEAR-lo, and (d) SPEAR-med compared with (a) observed drift from OSISAF. Observed and predicted JAS SIC (%) is plotted in colors.